

anti C4d – Literature List

Deceased donor kidney transplantation across donor-specific antibody barriers: predictors of antibody-mediated rejection

Elisabeth Schwaiger, Farsad Eskandary, Nicolas Kozakowski, Gregor Bond, Zeljko Kikic, Daniel Yoo, Susanne Rasoul-Rockenschaub, Rainer Oberbauer, and Georg A. Böhning

Nephrol. Dial. Transplant., Aug 2016; 31: 1342-1351.

Value of Donor-Specific Anti-HLA Antibody Monitoring and Characterization for Risk Stratification of Kidney Allograft Loss

Denis Viglietti, Alexandre Loupy, Dewi Vernerey, Carol Bentlejewski, Clément Gosset, Olivier Aubert, Jean-Paul Duong van Huyen, Xavier Jouven, Christophe Legendre, Denis Glotz, Adriana Zeevi, and Carmen Lefaucheur

J. Am. Soc. Nephrol., Aug 2016; 10.1681/ASN.2016030368.

Ig-like transcript 4 as a cellular receptor for soluble complement fragment C4d

Johannes Hofer, Florian Forster, David E. Isenman, Markus Wahrmann, Judith Leitner, Markus A. Hözl, Johannes J. Kovarik, Hannes Stockinger, Georg A. Böhning, Peter Steinberger, and Gerhard J. Zlabinger

FASEB J., Apr 2016; 30: 1492-1503.

IgG Donor-Specific Anti-Human HLA Antibody Subclasses and Kidney Allograft Antibody-Mediated Injury

Carmen Lefaucheur, Denis Viglietti, Carol Bentlejewski, Jean-Paul Duong van Huyen, Dewi Vernerey, Olivier Aubert, Jérôme Verine, Xavier Jouven, Christophe Legendre, Denis Glotz, Alexandre Loupy, and Adriana Zeevi

J. Am. Soc. Nephrol., Jan 2016; 27: 293-304.

The 20S proteasome core, active within apoptotic exosome-like vesicles, induces autoantibody production and accelerates rejection

Mélanie Dieudé, Christina Bell, Julie Turgeon, Deborah Beillevaire, Luc Pomerleau, Bing Yang, Katia Hamelin, Shijie Qi, Nicolas Pallet, Chanel Béland, Wahiba Dhahri, Jean-François Cailhier, Matthieu Rousseau, Anne-Claire Duchez, Tania Lévesque, Arthur Lau, Christiane Rondeau, Diane Gingras, Danie Muruve, Alain Rivard, Hélène Cardinal, Claude Perreault, Michel Desjardins, Éric Boilard, Pierre Thibault, and Marie-Josée Hébert

Science Translational Medicine, Dec 2015; 7: 318ra200.

Complement Factor C4d Is a Common Denominator in Thrombotic Microangiopathy

Jamie S. Chua, Hans J. Baelde, Malu Zandbergen, Suzanne Wilhelmus, Leendert A. van Es, Johan W. de Fijter, Jan A. Bruijn, Ingeborg M. Bajema, and Danielle Cohen

J. Am. Soc. Nephrol., Sep 2015; 26: 2239-2247.

Determinants and Outcomes of Accelerated Arteriosclerosis: Major Impact of Circulating Antibodies

Alexandre Loupy, Dewi Vernerey, Denis Viglietti, Olivier Aubert, Jean-Paul Duong van Huyen, Jean-Philippe Empana, Patrick Bruneval, Denis Glotz, Christophe Legendre, Xavier Jouven, and Carmen Lefaucheur
Circ. Res., Aug 2015; 117: 470-482.

Classical Complement Pathway Activation in the Kidneys of Women With Preeclampsia

Marlies Penning, Jamie S. Chua, Cees van Kooten, Malu Zandbergen, Aletta Buurma, Joke Schutte, Jan Anthonie Bruijn, Eliyahu V. Khankin, Kitty Bloemenkamp, S. Ananth Karumanchi, and Hans Baelde
Hypertension, Jul 2015; 66: 117-125.

Subclinical Rejection Phenotypes at 1 Year Post-Transplant and Outcome of Kidney Allografts

Alexandre Loupy, Dewi Vernerey, Claire Tinel, Olivier Aubert, Jean-Paul Duong van Huyen, Marion Rabant, Jérôme Verine, Dominique Nochy, Jean-Philippe Empana, Frank Martinez, Denis Glotz, Xavier Jouven, Christophe Legendre, and Carmen Lefaucheur

J. Am. Soc. Nephrol., Jul 2015; 26: 1721-1731.

Molecular Microscope Strategy to Improve Risk Stratification in Early Antibody-Mediated Kidney Allograft Rejection

Alexandre Loupy, Carmen Lefaucheur, Dewi Vernerey, Jessica Chang, Luis G. Hidalgo, Thibaut Beuscart, Jerome Verine, Olivier Aubert, Sébastien Dubleumortier, Jean-Paul Duong van Huyen, Xavier Jouven, Denis Glotz, Christophe Legendre, and Philip F. Halloran

J. Am. Soc. Nephrol., Oct 2014; 25: 2267-2277.

Stable long-term mixed chimerism achieved in a canine model of allogeneic in utero hematopoietic cell transplantation

Jesse D. Vrecenak, Erik G. Pearson, Matthew T. Santore, Carlyn A. Todorow, Haiying Li, Antoneta Radu, Tricia Bhatti, William H. Peranteau, Mark P. Johnson, and Alan W. Flake

Blood, Sep 2014; 124: 1987-1995.

Association of C4d Deposition with Clinical Outcomes in IgA Nephropathy

Mario Espinosa, Rosa Ortega, Marina Sánchez, Alfons Segarra, María Teresa Salcedo, Fayna González, Rafael Camacho, Miguel Angel Valdivia, Rocio Cabrera, Katia López, Fernando Pinedo, Eduardo Gutierrez, Alfonso Valera, Miryam Leon, María Angeles Cobo, Rosa Rodriguez, Jose Ballarín, Yolanda Arce, Beatriz García, María Dolores Muñoz, Manuel Praga for the Spanish Group for the Study of Glomerular Diseases (GLOSEN)

Clin. J. Am. Soc. Nephrol., May 2014; 9: 897-904.

Expression of hypoxia-inducible factor-1α (HIF-1α) in infiltrating inflammatory cells is associated with chronic allograft dysfunction and predicts long-term graft survival

Tung-Min Yu, Mei-Chin Wen, Chi-Yuan Li, Chi-Hung Cheng, Ming-Ju Wu, Cheng-Hsu Chen, and Kuo-Hsiung Shu

Nephrol. Dial. Transplant., Mar 2013; 28: 659-670.

Preeclampsia Is Characterized by Placental Complement Dysregulation

Aletta Buurma, Danielle Cohen, Kimberley Veraar, Dorrit Schonkeren, Frans H. Claas, Jan A. Bruijn, Kitty W. Bloemenkamp, and Hans J. Baelde

Hypertension, Nov 2012; 60: 1332-1337.

Solid phase detection of C4d-fixing HLA antibodies to predict rejection in high immunological risk kidney transplant recipients.

G Bartel, M Wahrmann, E Schwaiger, Z Kikic, C Winzer, WH Horl, F Muhlbacher, M Hoke, GJ Zlabinger, H Regele, and GA Bohmig

Transpl Int, Nov 2012

Long-Term Outcomes of Pre-emptive Valganciclovir Compared with Valacyclovir Prophylaxis for Prevention of Cytomegalovirus in Renal Transplantation

Tomas Reischig, Petra Hribova, Pavel Jindra, Ondrej Hes, Mirko Bouda, Vladislav Treska, and Ondrej Viklicky

J. Am. Soc. Nephrol., Sep 2012; 23: 1588-1597.

Anti-A/B antibody depletion by semiselective versus ABO blood group-specific immunoabsorption

Markus Wahrmann, Martin Schiemann, Lena Marinova, Günther F. Körmöczi, Kurt Derfler, Thomas Fehr, Georg Stussi, and Georg A. Böhmig

Nephrol. Dial. Transplant., May 2012; 27: 2122-2129.

Comparison between bortezomib and rituximab in the treatment of antibody-mediated renal allograft rejection

Johannes Waisser, Klemens Budde, Manuela Schütz, Lutz Liefeldt, Birgit Rudolph, Constanze Schönemann, Hans-H. Neumayer, and Nils Lachmann

Nephrol. Dial. Transplant., Mar 2012; 27: 1246-1251.

C4d Staining In Renal Allograft Biopsies with Early Acute Rejection and Subsequent Clinical Outcome

Johanna M Botermans, Hanneke de Kort, Michael Eikmans, Klaas Koop, Hans J. Baelde, Marko J.K. Mallat, Kim Zuidwijk, Cees van Kooten, Emile de Heer, Natascha N.T. Goemaere, Frans H.J. Claas, Jan A. Bruijn, Johan W de Fijter, Ingeborg M. Bajema, and Marian C. van Groningen

Clin. J. Am. Soc. Nephrol., May 2011; 6: 1207-1213.

Donor-Specific Antibodies Accelerate Arteriosclerosis after Kidney Transplantation

Gary S. Hill, Dominique Nochy, Patrick Bruneval, J. P. Duong van Huyen, Denis Glotz, Caroline Suberbielle, Julien Zuber, Dany Anglicheau, Jean-Philippe Empana,

Christophe Legendre, and Alexandre Loupy
J. Am. Soc. Nephrol., May 2011; 22: 975-983.

C4d staining in post-reperfusion renal biopsy is not useful for the early detection of antibody-mediated rejection when CDC crossmatching is negative

Elias David-Neto, Daisa S. R. David, Giordano F. Ginani, Helcio Rodrigues, Patricia S. Souza, Maria Cristina R. Castro, Hideki Kanashiro, Fernando Saito, Renato Falci, Jr, Ioannis M. Antonopoulos, Afonso Celso Piovesan, and William C. Nahas
Nephrol. Dial. Transplant., Apr 2011; 26: 1388-1392.

Prevalence and qualitative properties of circulating anti-human leukocyte antigen alloantibodies after pregnancy: no association with unexplained recurrent miscarriage.

Bartel G, Walch K, Wahrmann M, Pils S, Küssel L, Polterauer S, Tempfer C, Böhmig GA.

Hum Immunol. 2011 Feb;72(2):187-92. Epub 2010 Nov 12.

Significance of peritubular capillary, glomerular, and arteriolar C4d staining patterns in paraffin sections of early kidney transplant biopsies.

Z Kikic, H Regele, V Nordmeyer, M Wahrmann, J Kletzmayr, G Bartel, and GA Bohmig

Transplantation, Feb 2011; 91(4): 440-446.

Immunohistochemical characterization of glomerular and tubulointerstitial infiltrates in renal transplant patients with chronic allograft dysfunction

Chiara Divella, Michele Rossini, Antonia Loverre, Antonio Schena, Annamaria Maiorano, Vincenzo Gesualdo, Gianluigi Zaza, Giuseppe Grandaliano, and Francesco Paolo Schena

Nephrol. Dial. Transplant., Dec 2010; 25: 4071-4077.

Association between anti- β 2 glycoprotein I antibodies and renal glomerular C4d deposition in lupus nephritis patients with glomerular microthrombosis: a prospective study of 155 cases

Y. Shen, X-W. Chen, C-Y. Sun, M. Dai, Y-C. Yan, and C-D. Yang

Lupus, Sep 2010; 19: 1195-1203.

C4d-fixing capability of low-level donor-specific HLA antibodies is not predictive for early antibody-mediated rejection.

G Honger, M Wahrmann, P Amico, H Hopfer, GA Bohmig, and S Schaub

Transplantation, Jun 2010; 89(12): 1471-1475.

Effect of the proteasome inhibitor bortezomib on humoral immunity in two presensitized renal transplant candidates.

Wahrmann M, Haidinger M, Körmöczí GF, Weichhart T, Säemann MD, Geyeregger R, Kikić Z, Prikoszovich T, Drach J, Böhmig GA.

Transplantation. 2010 Jun 15;89(11):1385-1390.

The histological development of acute antibody-mediated rejection in HLA antibody-incompatible renal transplantation

Rob Higgins, Daniel Zehnder, Klaus Chen, David Lowe3 Joanna McKinnell, For T. Lam, Habib Kashi, Lam Chin Tan, Chris Imray, Simon Fletcher, Nithya Krishnan, Rizwan Hamer and David Briggs

Nephrol Dial Transplant (2010) 25: 1306-1312

Approach to a lung transplant biopsy

D M Hwang and S A Yousem

J. Clin. Pathol., Jan 2010; 63: 38-46.

Clinical relevance of preformed C4d-fixing and non-C4d-fixing HLA single antigen reactivity in renal allograft recipients.

M Wahrmann, G Bartel, M Exner, H Regele, GF Kormoczi, GF Fischer, and GA Böhmig

Transpl Int, Oct 2009; 22(10): 982-989.

Monocytes/macrophages in kidney allograft intimal arteritis: no association with markers of humoral rejection or with inferior outcome

Nicolas Kozakowski, Georg A. Böhmig, Markus Exner, Afschin Soleiman, Nicole Huttary, Katalin Nagy-Bojarszky, Rupert C. Ecker, Zeljko Kikic, and Heinz Regele
Nephrol. Dial. Transplant., Jun 2009; 24: 1979-1986.

Mesangial C4d deposition: a new prognostic factor in IgA nephropathy

Mario Espinosa, Rosa Ortega, Jose Manuel Gómez-Carrasco, Fernando López-Rubio, María López-Andreu, María Ovidia López-Oliva, and Pedro Aljama

Nephrol. Dial. Transplant., Mar 2009; 24: 886-891.

The significance of renal C4d staining in patients with BK viruria, viremia, and nephropathy.

Ibrahim Batal, Hanady Zainah, Sean Stockhausen, Amit Basu, Henkie Tan, Ron Shapiro, Adriana Zeevi, Alin Girnita and Parmjeet Randhawa

Modern Pathology, 2009; 22:1468-1476

Presence of FoxP3⁺ Regulatory T Cells Predicts Outcome of Subclinical Rejection of Renal Allografts

Oriol Bestard, Josep M. Cruzado, Inés Rama, Joan Torras, Montse Gomà, Daniel Serón, Francesc Moreso, Salvador Gil-Vernet, and Josep M. Grinyó

J. Am. Soc. Nephrol., Oct 2008; 19: 2020-2026.

Anti-glutathione S-transferase T1 antibody-mediated rejection in C4d-positive renal allograft recipients

Isabel Aguilera, Antonia Alvarez-Marquez, Miguel Angel Gentil, Jorge Fernandez-Alonso, Julia Fijo, Carmen Saez, Ingeborg Wichmann and Antonio Nuñez-Roldan
Nephrol Dial Transplant (2008) 23: 2393–2398

Antibodies, isotypes and complement in allograft rejection.

GA Bohmig, G Bartel, and M Wahrmann

Curr Opin Organ Transplant, Aug 2008; 13(4): 411-418.

Hydrolysis of Coagulation Factors by Circulating IgG Is Associated with a Reduced Risk for Chronic Allograft Nephropathy in Renal Transplanted Patients

Bharath Wootla, Antonino Nicoletti, Natacha Patey, Jordan D. Dimitrov, Christophe Legendre, Olivier D. Christophe, Alain Friboulet, Srinivas V. Kaveri, Sébastien Lacroix-Desmazes, and Olivier Thaunat

J. Immunol., Jun 2008; 180: 8455-8460.

The role of C4d immunostaining in the evaluation of the causes of renal allograft dysfunction

Prabhat Ranjan, Ritambhra Nada, Vivekanand Jha, Vinay Sakuja, and Kusum Joshi
Nephrol. Dial. Transplant., May 2008; 23: 1735-1741.

In vitro detection of C4d-fixing HLA alloantibodies: associations with capillary C4d deposition in kidney allografts.

G Bartel, M Wahrmann, M Exner, H Regele, N Huttary, M Schillinger, GF Kormoczi, WH Horl, and GA Bohmig

Am J Transplant, Jan 2008; 8(1): 41-49.

Achieving Donor-Specific Hyporesponsiveness Is Associated with FOXP3⁺ Regulatory T Cell Recruitment in Human Renal Allograft Infiltrates

Oriol Bestard, Josep M. Cruzado, Mariona Mestre, Anna Caldés, Jordi Bas, Marta Carrera, Joan Torras, Inés Rama, Francesc Moreso, Daniel Serón, and Josep M. Grinyó

J. Immunol., Oct 2007; 179: 4901-4909.

Demonstration of secretory IgA in kidneys of patients with IgA nephropathy

Beatrijs D. Oortwijn, Maria Pia Rastaldi, Anja Roos, Deborah Mattinzoli, Mohamed R. Daha and Cees van Kooten

Nephrology Dialysis Transplantation, 2007

C4d staining of renal allograft biopsies: a comparative analysis of different staining techniques.

Christian A. Seemayer, Ariana Gaspert, Volker Nickeleit and Michael J. Mihatsch

Nephrology Dialysis Transplantation 2007 22(2): 568-576

Determinants of the complement-fixing ability of recipient presensitization against HLA antigens.

G Bartel, M Wahrmann, M Exner, H Regele, M Schillinger, WH Horl, and GA Bohmig
Transplantation, Mar 2007; 83(6): 727-733.

Immunoabsorption in severe C4d-positive acute kidney allograft rejection: a randomized controlled trial.

GA Bohmig, M Wahrmann, H Regele, M Exner, B Robl, K Derfler, T Soliman, P Bauer, M Mullner, and W Druml

Am J Transplant, Jan 2007; 7(1): 117-121.

Comparison of C4d Immunostaining Methods in Renal Allograft Biopsies

Megan L. Troxell, Lauren A. Weintraub, John P. Higgins, and Neeraja Kambham

Clin. J. Am. Soc. Nephrol., May 2006; 1: 583-591.

Immunohistochemical evidence of activated lectin pathway in kidney allografts with peritubular capillary C4d deposition.

Naofumi Imai, Shinichi Nishi, Bassam Alchi, Mitsuhiro Ueno, Sachiko Fukase, Masaaki Arakawa, Kazuhide Saito, Kota Takahashi and Fumitake Gejyo

Nephrology Dialysis Transplantation 2006 21(9): 2589-2595

Rescue therapy with tacrolimus and mycophenolate mofetil does not prevent deterioration of graft function in C4d-positive chronic allograft nephropathy.

C Schwarz, H Regele, N Huttary, M Wahrmann, M Exner, K Nagy-Bojarsky, J Kletzmayr, WH Horl, and GA Bohmig

Wien Klin Wochenschr, Jul 2006; 118(13-14): 397-404.

Glomerular Activation of the Lectin Pathway of Complement in IgA Nephropathy Is Associated with More Severe Renal Disease

Anja Roos, Maria Pia Rastaldi, Novella Calvaresi, Beatrijs D. Oortwijn, Nicole Schlagwein, Danielle J. van Gijlswijk-Janssen, Gregory L. Stahl, Misao Matsushita, Teizo Fujita, Cees van Kooten, and Mohamed R. Daha

J. Am. Soc. Nephrol., Jun 2006; 17: 1724-1734.

Pivotal role of complement-fixing HLA alloantibodies in presensitized kidney allograft recipients.

M Wahrmann, M Exner, M Schillinger, B Haidbauer, H Regele, GF Kormoczi, WH Horl, and GA Bohmig

Am J Transplant, May 2006; 6(5 Pt 1): 1033-1041.

Comparison of C4d Immunostaining Methods in Renal Allograft Biopsies

Megan L. Troxell, Lauren A. Weintraub, John P. Higgins, and Neeraja Kambham

Clin. J. Am. Soc. Nephrol., May 2006; 1: 583-591.

Severe early acute humoral rejection resulting in allograft loss in a renal transplant recipient with Campath-1H induction therapy

Prue Hill, Elena Gagliardini, Piero Ruggenenti, and Giuseppe Remuzzi

Nephrol. Dial. Transplant., Aug 2005; 20: 1741-1744.

[C4d]FlowPRA screening--a specific assay for selective detection of complement-activating anti-HLA alloantibodies.

M Wahrmann, M Exner, B Haidbauer, M Schillinger, H Regele, G Kormoczi, and GA

Bohmig
Hum Immunol, May 2005; 66(5): 526-534.

Risk factors for capillary C4d deposition in kidney allografts: evaluation of a large study cohort.

M Lorenz, H Regele, M Schillinger, M Exner, S Rasoul-Rockenschaub, M Wahrmann, J Kletzmayr, G Silberhuber, WH Hörl, and GA Bohmig
Transplantation, Aug 2004; 78(3): 447-452.

Description of B Lymphocytes and Plasma Cells, Complement, and Chemokines / Receptors in Acute Liver Allograft Rejection.

Manfred G Krukemeyer, et al,
Transplantation 2004, 78(1)

Flow cytometry based detection of HLA alloantibody mediated classical complement activation.

M Wahrmann, M Exner, H Regele, K Derfler, GF Kormoczi, K Lhotta, GJ Zlabinger, and GA Bohmig
J Immunol Methods, Apr 2003; 275(1-2): 149-160.

Kidney Transplants, antibodies and rejections: is C4d a magic marker?

Volker Nickeleit and Michael J. Mihatsch
Nephrol. Dial Transplant 2003, 18: 2232-2239.

Antibody-mediated rejection criteria - an addition to the Banff 97 classification of renal allograft rejection.

Racusen LC, et al.
Am J Transplant 2003, 3: 708-714

Capillary Deposition of Complement Split Product C4d in Renal Allografts is Associated with Basement Membrane Injury in Peritubular and Glomerular Capillaries: A Contribution of Humoral Immunity to Chronic Allograft Rejection

Regele, Böhmig, Habicht, Gollowitzer, Schillinger, Rockenschaub, Watschinger, Kerjaschki, Exner
J Am Soc Nephrol 2002, 13: 2371-2380

Capillary C4d Deposition in Kidney Allografts: A Specific Marker of Alloantibody-Dependent Graft Injury.

Böhmig, Exner, Habicht, Schillinger, Lang, Kletzmayr, Säemann, Hörl, Watschiner, Regele

J Am Soc Nephrol 2002, 13: 1091-1099

C4d-Positive Acute Humoral Renal Allograft Rejection: Effective Treatment by Immunoadsorption.

Böhmig, Regele, Exner, Derhartunian, Kletzmayr, Säemann, Hörl, Druml, Watschiner
J Am Soc Nephrol 2001, 12: 2482-2489

Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection.

Regele, Exner, Watschinger, Wenter, Wahrmann, Österreicher, Säemann, Mersich, Hörl, Zlabiner, Böhming

Nephrol Dial Transplant 2001, 16: 2058-2066

C4d deposits in renal allografts are associated with inferior graft outcome.

GA Bohmig, M Exner, B Watschinger, C Wenter, M Wahrmann, C Österreicher, MD Saemann, N Mersich, WH Horl, GJ Zlabinger, and H Regele

Transplant Proc, Feb 2001; 33(1-2): 1151-1152.